

$^{13}\text{CO}(J = 1 - 0)$ DEPRESSION IN LUMINOUS STARBURST MERGERS

Yoshiaki Taniguchi and Youichi Ohyama

Astronomical Institute, Tohoku University, Aoba, Sendai 980-8578, Japan

ABSTRACT

It is known that the class of luminous starburst galaxies tends to have higher $R = ^{12}\text{CO}(J=1-0)/^{13}\text{CO}(J=1-0)$ integrated line intensity ratios ($R > 20$) than normal spiral galaxies ($R \sim 10$). Since most previous studies investigated only R , it remains uncertain whether the luminous starburst galaxies are overabundant in ^{12}CO or underabundant in ^{13}CO . Here we propose a new observational test to examine this problem. Our new test is to compare far-infrared luminosities [$L(\text{FIR})$] with those of ^{12}CO and ^{13}CO [$L(^{12}\text{CO})$ and $L(^{13}\text{CO})$, respectively]. It is shown that there is a very tight correlation between $L(^{12}\text{CO})$ and $L(\text{FIR})$, as found in many previous studies. However, we find that the ^{13}CO luminosities of the high- R galaxies are lower by a factor of three on the average than those expected from the correlation for the remaining galaxies with ordinary R values. Therefore, we conclude that the observed high R values for the luminous starburst galaxies are attributed to their lower ^{13}CO line intensities.

Subject headings: galaxies: emission lines - galaxies: starburst - interstellar: molecules

1. INTRODUCTION

It is known that the class of luminous starburst galaxies with $L(\text{FIR}) \gtrsim 10^{11} L_{\odot}$ tends to have higher $R = ^{12}\text{CO}(J=1-0)/^{13}\text{CO}(J=1-0)$ integrated line intensity ratios than normal spiral galaxies (Aalto et al. 1991, 1995, 1997; Casoli et al. 1991; Casoli, Dupraz, & Combes 1992a, 1992b; Hurt & Turner 1991; Turner & Hurt 1992; Garay, Mardones, & Mirabel 1993; Henkel & Mauersberger 1993; Henkel et al. 1998). It was often considered that the higher R values observed in these galaxies are attributed to the inflow of disk gas with high $^{12}\text{C}/^{13}\text{C}$ abundance ratios, possibly combined with a ^{12}C enhancement caused by nucleosynthesis in massive stars (e.g., Henkel et al. 1998). However, since most previous studies investigated only the ratios, it is uncertain whether or not these luminous starburst galaxies are overluminous in ^{12}CO or underluminous in ^{13}CO (Casoli et al. 1992a). In this Letter, comparing the luminosities of $^{12}\text{CO}(J=1-0)$, $^{13}\text{CO}(J=1-0)$ (hereafter ^{12}CO and ^{13}CO , respectively), and FIR emission for a sample of starburst and normal galaxies for which all the data are available in literature, we show that the observed higher R values are mainly attributed to a lower intensity in ^{13}CO with respect to ^{12}CO . We discuss possible physical mechanisms which can explain the $^{13}\text{CO}(J=1-0)$ depression in luminous starburst galaxies.

2. DATA AND RESULTS

We have compiled ^{12}CO and ^{13}CO intensities from the literature (Wiklind & Henkel 1990; Aalto et al. 1991, 1995; Becker & Freudling 1991; Sage & Isbell 1991; Casoli et al. 1992b; Garay et al. 1993; Henkel, Whiteoak, & Mauersberger 1994). Our sample consists of 51 galaxies including ultraluminous infrared galaxies such as Arp 220. These data are used to estimate the CO luminosities; $L(\text{CO})$ is defined

as $L(\text{CO}) = A \times I(\text{CO}) \text{ K km s}^{-1} \text{ pc}^2$ where A is the observed area in units of pc^2 and $I(\text{CO}) = \int T_{\text{A}}^* \eta^{-1} dv \text{ K km s}^{-1}$ where T_{A}^* is the observed antenna temperature corrected for atmospheric extinction and η is the main beam efficiency.

The FIR data are compiled from the IRAS Faint Source Catalog (Moshir et al. 1992). The FIR luminosities are estimated using $L(\text{FIR}) = 4\pi D^2 1.26 \times 10^{-11} [2.58 \times S(60) + S(100)] \text{ (ergs s}^{-1})$ where $S(60)$ and $S(100)$ are the IRAS 60 μm and 100 μm fluxes in units of Jy and D is the distance of galaxies in units of cm (Helou, Soifer, & Rowan-Robinson 1985). Distances of nearby galaxies are taken from the Nearby Galaxies Catalog (Tully 1988); distances of other galaxies are estimated using a Hubble constant $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ with V_{GSR} (radial velocity corrected to the Galactic Center: de Vaucouleurs et al. 1991). The compiled data are given in Table 1. Though our sample is not statistically complete, it is the largest sample compiled so far.

As shown in Table 1, there are several galaxies which have $R \geq 20$. Hereafter we refer to these sources as high- R galaxies (NGC 1614, NGC 3256, NGC 4194, NGC 6240, Arp 220, and Arp 299). ESO 541-IG 23 [= Arp 236, $R \geq 17$ (lower limit)] and possibly IRAS 18293–3413 ($R \simeq 26$: Garay et al. 1993; $R \simeq 17$: Aalto et al. 1995) are also high- R galaxies. In total, there are eight high- R galaxies in our sample. Except for the uncertain case of IRAS 18293–3413, all these high- R galaxies are mergers. For the remaining 43 galaxies, we obtain an average ratio, $\langle R \rangle = 11.3 \pm 3.3$. The values are consistent with previous estimates of 11 ± 3 (Aalto et al. 1991) and 9.3 ± 3.6 (Sage & Isbell 1991) for normal galaxies.

First, we examine whether or not there is a beam size effect on R because the CO line observations did not cover the whole area of the individual galaxies. Ratios of beam size (D_{beam} = half power beam width) to optical diameter at 25 mag arcsec $^{-2}$ (D_{25} : de Vaucouleurs et al. 1991) are given in the third column of Table 1. In Figure 1, D_{beam}/D_{25} is plotted as a function of R . We find no clear correlation, indicating that the difference in R cannot be attributed to the beam-size effect.

In Figure 2, we compare $L(^{12}\text{CO})$ with $L(^{13}\text{CO})$. A tight correlation is found for galaxies with $L(^{12}\text{CO}) < 10^{8.75} \text{ K km s}^{-1} \text{ pc}^2$. This almost linear correlation can be expressed as $\log L(^{13}\text{CO}) = (0.96 \pm 0.03) \times \log L(^{12}\text{CO}) + (-0.68 \pm 0.27)$. However, more luminous galaxies do not follow the same correlation, i.e., some of the luminous galaxies have lower ^{13}CO luminosities with respect to ^{12}CO .

In Figure 3, we compare $L(\text{FIR})$ with both $L(^{12}\text{CO})$ and $L(^{13}\text{CO})$. It is shown that $L(^{12}\text{CO})$ is well correlated with $L(\text{FIR})$; its correlation coefficient is 0.94. The average $L(^{12}\text{CO})/L(\text{FIR})$ for the high- R galaxies, 0.013 ± 0.007 , is similar to that for the normal ones, 0.023 ± 0.013 . This correlation has been noted in many previous studies (e.g., Young & Scoville 1991 and references therein). In fact, Figure 3 shows that the correlation is established within a factor of three dispersion except for one galaxy (NGC 55) which is a nearby Magellanic-type irregular galaxy (Becker & Freudling 1991). On the other hand, the correlation between $L(\text{FIR})$ and $L(^{13}\text{CO})$ appears to be significantly poorer (the correlation coefficient is 0.86) than that between $L(\text{FIR})$ and $L(^{12}\text{CO})$. The correlation is poorer because the high- R luminous starburst galaxies have lower ^{13}CO luminosities than expected for galaxies with ordinary R values; note that the average value of $L(^{13}\text{CO})/L(\text{FIR})$ is 0.00051 ± 0.00034 for the high- R galaxies while that for the normal galaxies is 0.0023 ± 0.0017 . Thus, we conclude that the higher R in the luminous starburst mergers is mostly due not to enhanced ^{12}CO emission but to depressed ^{13}CO line intensities. An average R value for the high- R galaxies is $\simeq 30.7 \pm 11.5$. Since lower-limit data are used in this estimate, we obtain a ^{13}CO depression factor $\gtrsim 3$, given $\langle R \rangle \simeq 11$ for the normal galaxies. Any physical mechanism must explain the intrinsic weakness of $^{13}\text{CO}(J = 1 - 0)$ emission by a factor of $\gtrsim 3$ in the luminous starburst mergers.

3. DISCUSSION

Here we investigate possible mechanisms that can explain the intrinsic weakness of $^{13}\text{CO}(J = 1 - 0)$ emission by a factor of $\gtrsim 3$ in the luminous starburst mergers. High R values may be caused by (see also Sakamoto et al. 1997): 1) low gas densities, 2) high gas kinetic temperatures, 3) large nuclear velocity gradients that keep CO column densities small at a given velocity, 4) small CO/H₂ abundance ratios, 5) more efficient excitation of ^{13}CO molecules toward upper levels (e.g., $J \geq 2$) with respect to ^{12}CO ones, or 6) large $^{12}\text{CO}/^{13}\text{CO}$ abundance ratios. The first four mechanisms lead to lower opacity of the CO lines. The first mechanism also introduces non-LTE excitation, resulting in high R values. Since it is known that the luminous starburst mergers have a huge amount of dense gas ($\sim 10^{10} M_{\odot}$; Solomon et al. 1992; Scoville, Yun, & Bryant 1997), the first idea (lower gas densities) seems to be unlikely. Starburst galaxies often have both high gas kinetic temperatures and particularly large velocity gradients (Turner, Martin, & Ho 1990; Aalto et al. 1991, 1995; Devereux et al. 1994). However, if these mechanisms were responsible for the higher R values, we would find a ^{12}CO excess with respect to ^{13}CO . Therefore the second and third idea are also ruled out. The smaller CO/H₂ abundance ratio (i.e., lower metallicity) seems also to be unlikely for the luminous starburst mergers because the enhanced star-formation activity occurs in their nuclear regions where metal abundances are generally high (e.g., Storch-Bergmann et al. 1996). It is known that $^{12}\text{CO}(J=3-2)/^{12}\text{CO}(J=1-0)$ ratios of starburst galaxies are often found to be higher than in normal galaxies (e.g., Devereux et al. 1994 and references therein). If ^{13}CO molecules in starburst mergers were excited more efficiently toward upper levels with respect to ^{12}CO , we would observe high- R values in higher rotational transitions. However, there is no such evidence (Taniguchi, Ohyama, & Sanders 1998). Therefore, the remaining possibility is that the $^{12}\text{CO}/^{13}\text{CO}$ abundance ratios are systematically higher than those in normal galaxies. According to our result (Fig. 3), this should be achieved by a ^{13}CO underabundance with respect to ^{12}CO in the starburst mergers.

We would like to thank Dave Sanders, Naomasa Nakai, Toshihiro Handa, Sumio Ishizuki, Naomi Hirano, and Satoki Matsushita for useful comments. We would also like to thank the referee, Christian Henkel, for his many useful comments and suggestions which improved this paper significantly. Y.O. was supported by the Grant-in-Aid for JSPS Fellows by the Ministry of Education, Science, Sports and Culture. This work was supported in part by the Ministry of Education, Science, Sports and Culture in Japan under Grant Nos. 07055044, 10044052, and 10304013.

REFERENCES

- Aalto, S., Black, J. H., Johansson, L. E. B., & Booth, R. S. 1991, *A&A*, 249, 323
- Aalto, S., Booth, R. S., Black, J. H., & Johansson, L. E. B. 1995, *A&A*, 300, 369
- Aalto, S., Radford, S. J. E., Scoville, N. Z., & Sargent, A. I. 1997, *ApJ*, 475, L107
- Becker, R., & Freudling, W. 1991, *A&A*, 251, 454
- Casoli, F., Dupraz, C., & Combes, F. 1992a, *A&A*, 264, 49
- Casoli, F., Dupraz, C., & Combes, F. 1992b, *A&A*, 264, 55
- Casoli, F., Dupraz, C., Combes, F., & Kazès, I. 1991, *A&A*, 251, 1
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G. Jr., Buta, R. J., Paturel, G., & Fouqué, P. 1991, *Third Reference Catalogue of Bright Galaxies* (Springer-Verlag)

- Devereux, N., Taniguchi, Y., Sanders, D. B., Nakai, N., & Young, J. S. 1994, *AJ*, 107, 2006
- Freeman, K. C., Karlsson, B., Lynga, G., Burrell, J. F., van Woerden, H., & Goss, R. 1977, *A&A*, 55, 445
- Garay, G., Mardones, D., & Mirabel, I. F. 1993, *A&A*, 277, 405
- Helou, G. I., Soifer, B. T., & Rowan-Robinson, M. 1985, *ApJ*, 298, L7
- Henkel, C., Chin, Y.-N., Mauersberger, R., & Whiteoak, J. B. 1998, *A&A*, 329, 443
- Henkel, C., & Mauersberger, R. 1993, *A&A*, 274, 730
- Henkel, C., Whiteoak, J. B., & Mauersberger, R. 1994, *A&A*, 284, 17
- Hurt, R. L., & Turner, J. L. 1991, *ApJ*, 377, 434
- Lonsdale, C. J., Helou, G., Good, J. C., & Rice, W. 1989, *Cataloged Galaxies and Quasars Observed in the IRAS Survey, Version 2*, JPL D-1932 (Pasadena: JPL)
- Moshir, M., et al. 1992, *Explanatory Supplement to the IRAS Faint Source Survey, Version 2*, JPL-D-10015 8/92 (Pasadena: JPL)
- Rice, W., Lonsdale, C. J., Soifer, B. T., Neugebauer, G., Kopan, E. L., Lloyd, L. A., de Jong, T., & Habing, H. J. 1988, *ApJS*, 68, 91
- Sage, L. J., & Isbell, D. W. 1991, *A&A*, 247, 320
- Sakamoto, S., Handa, T., Sofue, Y., Honma, M., & Sorai, K. 1997, *ApJ*, 475, 134
- Sanders, D. B., Egami, E., Lipari, S., Mirabel, I. F., & Soifer, B. T. 1995, *AJ*, 110, 1993
- Scoville, N. Z., Yun, M. S., & Bryant, P. M. 1997, *ApJ*, 484, 702
- Solomon, P. M., Downes, D., & Radford, S. J. E. 1992, *ApJ*, 387, L55
- Storchi-Bergmann, T., Rodriguez-Ardila, A., Schmitt, H. R., Wilson, A. S., & Baldwin, J. A. 1996, *ApJ*, 472, 83
- Strauss, M. A., Huchra, J. P., Davis, M., Yahil, A., Fisher, K. B., & Tonry, J. 1992, *ApJS*, 83, 29
- Taniguchi, Y., Ohya, Y., & Sanders, D. B. 1998, in preparation
- Tully, R. B. 1988, *Nearby Galaxies Catalog* (Cambridge University Press)
- Turner, J. L., & Hurt, R. L. 1992, *ApJ*, 384, 72
- Turner, J. L., Martin, R. N., & Ho, P. T. P. 1990, *ApJ*, 351, 418
- Wiklind, T., & Henkel, C. 1990, *A&A*, 227, 394
- Young, J. S., & Scoville, N. Z. 1991, *ARA&A*, 29, 581

Table 1. Properties of a sample of galaxies

Name	d (Mpc)	D_{beam} $/D_{25}$	$L(^{12}\text{CO})$ (K km s ⁻¹ pc ²)	$L(^{13}\text{CO})$ (K km s ⁻¹ pc ²)	R	$L(\text{FIR})$ ($L_{\odot}$)	Ref. ^a
N55	1	0.023	5.51	4.33	15.4	8.03	1
N134	19	0.084	8.29	7.14	14.0	10.16	2
N253	3	0.034	8.26	7.04	16.6	10.03	3
N404	2	0.106	5.75	4.79	9.1	7.33	4
N520	28	0.123	8.76	7.71	11.0	10.58	5
N628	10	0.091	7.57	6.75	6.6	9.64 ^b	3
N660	12	0.066	8.58	7.43	14.1	10.17	5
N828	73 ^c	0.191	9.73	8.70	10.7	11.03	5
...	...	0.127	9.61	8.44	14.9	...	6
N891	10	0.070	8.40	7.48	8.4	9.90	3
N986	23	0.184	8.75	7.74	10.2	10.35	5
N1058	9	0.315	7.13	6.07	11.5	8.70	3
N1614	62 ^c	0.544	9.11	<7.67	>27.4	11.25	5
N1808	11	0.111	8.46	7.25	16.5	10.22	5
N2146	17	0.091	8.82	7.74	12.0	10.78	5
N2276	37	0.195	8.66	7.75	8.0	10.50	2
N2369	44 ^c	0.202	9.39	8.22	15.0	10.82	2
...	...	0.211	9.33	8.28	11.3	...	7
N2903	6	0.075	7.79	6.81	9.5	9.48	3
N3034	5	0.085	8.67	7.47	15.9	10.62	3
N3079	20	0.069	8.94	7.91	10.8	10.52	5
N3256	37	0.189	9.57	8.01	35.6	11.28	5
...	...	0.193	9.54	8.07	29.2	...	1
...	...	0.096	8.95	7.40	35.5	...	6
...	...	0.197	9.53	8.12	25.9	...	7
N3627	7	0.104	7.94	6.77	14.8	9.57	3
N3628	8	0.064	8.34	7.32	10.5	9.73	3
N4038/9	25	0.086 ^d	8.94	7.82	13.3	10.64	5
N4194	39	0.201	8.83	7.09	54.4	10.69	6
N4414	10	0.262	8.27	7.47	6.3	9.70	3
N4736	4	0.085	7.29	6.37	8.3	9.33 ^b	3
N4826	4	0.055	7.22	6.51	5.1	9.03	5
...	...	0.095	7.50	6.77	5.3	...	3
N4945	5	0.036	8.61	7.45	14.4	10.22	8
...	...	0.036	8.63	7.45	15.1	...	5
N5033	19	0.051	8.27	7.32	8.9	10.04	5
N5055	7	0.044	7.65	6.86	6.2	9.57	5
N5218	39 ^c	0.302	9.13	8.16	9.3	10.28	5
N5457	5	0.074	8.53 ^e	7.69 ^e	6.9 ^e	9.74 ^b	3
N6215	21	0.335	8.21	7.13	12.1	10.33 ^f	2
N6221	19	0.202	8.60	7.35	18.0	10.49 ^f	2

Table 1—Continued

Name	d (Mpc)	D_{beam} $/D_{25}$	$L(^{12}\text{CO})$ (K km s $^{-1}$ pc 2)	$L(^{13}\text{CO})$ (K km s $^{-1}$ pc 2)	R	$L(\text{FIR})$ ($L_{\odot}$)	Ref. ^a
N6240	98 ^c	0.172	9.72	8.08	43.6	11.52	6
N6503	6	0.134	7.23	6.11	13.2	8.82	3
N6810	25	0.227	8.61	7.65	9.0	10.31	2
N6946	6	0.083	8.16	7.12	11.1	9.55	3
N7130	65 ^c	0.473	9.21	8.31	8.0	11.05	5
N7552	20	0.212	8.72	7.56	14.2	10.63	2
N7582	18	0.143	8.50	7.28	16.7	10.39	2
IC342	4	0.044	7.81	6.76	11.1	9.32	3
IC2554	17	1.136	8.09	6.97	13.0	9.95 ^f	2
U2855	20	0.126	8.52	7.44	11.9	10.40	2
Circinus	4 ^g	0.104	7.85	6.83	10.5	9.78 ^h	5
Maffei2	3	0.950	7.64	6.68	9.0	9.33 ^h	3
Arp220	74 ^c	0.363	9.58	<8.34	>17.3	11.91	5
...	...	0.242	9.72	<8.39	>21.8	...	6
Arp299	42 ^c	0.468	9.09	<7.80	>19.8	11.41	5
...	...	0.312	8.99	7.68	20.7	...	6
IRAS18293 ⁱ	73 ^j	2.389 ^k	9.79	8.56	16.8	11.46	2
...	...	2.500 ^k	9.86	8.44	25.9	...	7
IRAS22132 ⁱ	46 ^c	0.320	9.07	8.12	8.9	10.84	7
ESO541-IG23	81 ^c	1.604	9.59	<8.36	>17.0	11.36	2

^a1: Becker & Freudling (1991), 2: Aalto et al. (1995), 3: Sage & Isbell (1991), 4: Wiklind & Henkel (1990), 5: Aalto et al. (1991), 6: Casoli et al. (1992b), 7: Garay et al. (1993), 8: Henkel et al. (1994)

^bRice et al. (1988)

^cEstimated using a Hubble constant $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ with V_{GSR} given in de Vaucouleurs et al. (1991).

^dSum of D_{25} of both N4038 and N4039 is used.

^eData in all regions (nuclus and four disk regions) are taken together (see Sage & Isbell 1991).

^fSanders et al. (1995)

^gFreeman et al. (1977)

^hLonsdale et al. (1989)

ⁱIRAS18293 = IRAS 18293-3413 and IRAS22132 = IRAS 22132-3705

^jStrauss et al. (1992)

^kOptical diameter is taken from NED.




Fig. 1.— Diagram of D_{beam}/D_{25} vs. R . Objects with $R \gtrsim 20$ (see the main text) are shown by filled circles and labeled. Note that IRAS18293 = IRAS 18293–3413 and ESO541 = ESO 541–IG 23.




Fig. 2.— Diagram of $L(^{12}\text{CO})$ vs. $L(^{13}\text{CO})$. The symbols have the same meaning as those in figure 1. A solid line corresponds to the correlation for $R = \langle R \rangle$ where $\langle R \rangle$ is the mean R using the data of galaxies with $R < 20$ ($\langle R \rangle = 11.3$). The dotted lines corresponds to the correlation for $R = \langle R \rangle \pm \sigma(R)$ where $\sigma(R) = 3.3$.




Fig. 3.— Diagrams of R (top), $L(^{12}\text{CO})$ (middle), and $L(^{13}\text{CO})$ (bottom) against $L(\text{FIR})$. The symbols have the same meaning as those in Figure 1. Alphabets in the top panel mean; a: NGC 4194, b: NGC 6240, c: NGC 3256, d: NGC 3256, e: NGC 3256, f: NGC 1614, g: NGC 3256, h: IRAS 18293–3413, i: Arp 220, j: Arp 299, k: Arp 299, l: Arp 220, m: ESO 541–IG 23, and n: IRAS 18293–3413.